

SFU-HEP-04-96

Revised 07-96

String Theory In Curved Space-Time

K.S.Viswanathan¹

Department of Physics
Simon Fraser University
Burnaby,B.C., Canada V5A 1S6

and

R.Parthasarathy²

The Institute of Mathematical Sciences
C.P.T.Campus, Taramani Post
Madras 600 113, India

Abstract

Intrinsic and extrinsic geometric properties of string world sheets in curved space-time background are explored. In our formulation, the only dynamical degrees of freedom of the string are its immersion coordinates. Classical equation of motion and the space-time energy-momentum tensor of the string are obtained. The equations of motion for the extrinsic curvature action are second order for the scalar mean curvature of the world sheet. 1-loop divergent terms are calculated using the background field method. Asymptotic freedom of the extrinsic curvature coupling is established.

PACS number(s): 11.25.-w, 11.25.Pm

¹Electronic address: kviswana@sfu.ca

²Electronic address: sarathy@imsc.ernet.in

0.1 INTRODUCTION

String theory in curved space-times is an exciting subject and in fact has been investigated by many authors [see [1] for a recent review] as a framework to study the physics of gravitation in the context of string theory. In most investigations so far, the starting point for open or closed bosonic strings propagating in a D-dimensional space-time with metric $h_{\mu\nu}(X)$; ($0 \leq \mu, \nu \leq D-1$) is the action

$$S = \frac{1}{2\pi\alpha'} \int d\sigma d\tau \sqrt{g} g_{\alpha\beta}(\sigma, \tau) h_{\mu\nu}(X) \partial_\alpha X^\mu(\sigma, \tau) \partial_\beta X^\nu(\sigma, \tau). \quad (1)$$

Here $g_{\alpha\beta}(\sigma, \tau)$ ($\alpha, \beta = 1, 2$) is the metric on the world sheet. α' is the string tension. The metric $h_{\mu\nu}$ may be taken as either a fixed background or as dynamically generated by the string. The dynamical variables of the theory described by (1) are the immersion coordinates $X^\mu(\sigma, \tau)$ and the world sheet metric $g_{\alpha\beta}$. Let us recall that the above action is obtained from the Nambu-Goto (N-G) action

$$S = \frac{1}{2\pi\alpha'} \int \sqrt{g} d\sigma d\tau, \quad (2)$$

which in turn can be written as

$$S = \frac{1}{2\pi\alpha'} \int (\sqrt{g} + \lambda^{\alpha\beta} (\partial_\alpha X^\mu \partial_\beta X^\nu h_{\mu\nu} - g_{\alpha\beta})) d\sigma d\tau, \quad (3)$$

and presuming condensation of the Lagrange multiplier field $\lambda_{\alpha\beta}$ [2]

$$\langle \lambda_{\alpha\beta} \rangle = \text{const.} \sqrt{g} g_{\alpha\beta}. \quad (4)$$

The reasons for starting with the alternative (1) in string theory are well known. The main points being: the integration over X^μ is easily done and the full power of the theory of Riemann surfaces can be exploited when integrating over the space of all 2-metrics.

In investigating strings in a curved background, it will be shown in this paper that it is more convenient to work with the N-G action (2), where the only dynamical variables are $X^\mu(\sigma, \tau)$ while the metric of the world sheet is induced from its immersion in background geometry. In this scheme we do not have to contend with the Weyl symmetry of the 2-d metric. It will

also become evident below that, since the geometric properties of the world sheet are described in terms of both its first and second fundamental forms, it is natural to take into account, besides the usual N-G action, an action that involves the extrinsic geometry of the world sheet. 2-d surfaces that extremize the N-G action have $H^i = 0$, where $H^i (i = 1, \dots, D - 2)$ are the scalar mean curvatures of the surface. A much wider class of surfaces extremize the extrinsic curvature action, thereby, enabling one to investigate the role of a very general class of world sheets on the dynamics of strings. There exists a large body of work [3]-[7] on strings with extrinsic curvature action in Lorentzian space-time with the hope of describing QCD strings. The necessity of including such a term for QCD was emphasized in [3]. Extrinsic geometry dependent term in the action provides rigidity to the strings, while the N-G term only provides tension. Rigidity effects are important also in determining the shapes of biological membranes [8],[9]. It is thus natural to investigate the role of extrinsic geometry on the properties of strings in curved space-time. String theory has a dimensional parameter, i.e., the string tension. Rigid strings have a dimensionless coupling which in flat background is asymptotically free. We shall see that this coupling is asymptotically free in curved background as well.

Although we discuss primarily string world sheet in curved background, many of the results obtained here may be used in the context of the (3+1) canonical formulation of gravity. It may be recalled that in the (3+1) formulation of Arnowitt, Deser and Misner (ADM) [10],[11], the Cauchy problem of general relativity can be tackled by constructing a space-time $\tilde{M}$ out of a foliation of space-like hypersurfaces with time coordinate t as a parameter. In this formalism the quantities needed for the description of the 4-dimensional line element are the lapse function, which measures distance between nearby hypersurfaces, the shift vector that gives the relation between the spatial coordinate systems on different hypersurfaces and the 3-d space metric to enable one to measure distances in space-like, $t = \text{constant}$ slices. In 4-d space-time, Einstein's theory leads to three momentum constraints, the Hamiltonian constraint, and six equations of motion. The hyperbolic form of Einstein's theory can be obtained [12],[13] by imposition of a slicing condition on the space-time. One such condition involves taking $t = \text{constant}$ surfaces to be constant mean curvature hypersurfaces. Thus, for example, using harmonic Gauss map, one can solve the constraints in (2+1) gravity [14]-[16]. The considerations in this paper offer other possibilities for choosing

hypersurfaces that extremize the extrinsic curvature action. 2-d surfaces of prescribed mean curvature are studied as solutions of the Einstein constraint equations on closed manifolds in [17],[18].

Earlier studies in string theory in curved background, for example, in [19] - [23], concentrated on the classical equation of motion in order to exhibit its nonlinearity. While the authors in [19], [22] and [23] considered Nambu-Goto action in curved background, [20] and [21] (in the second paper) included extrinsic curvature dependent term in the action. In these studies the first variation of the action giving the classical equations of motion is examined for various choices of the background spacetime. Some of the classical aspects of strings in curved background discussed in this paper have also been considered recently by Capovilla and Guven [24]. They have considered the geometry of deformations of relativistic membranes. In particular they have derived the second variation of the NG action in a general curved space and that of the extrinsic curvature action in a background Minkowski space-time. In this paper we are interested in both the classical and the quantum aspects of strings in curved space. In this paper we calculate the second variation of both the NG and the extrinsic curvature actions in arbitrary curved space-time which is then used to calculate the renormalization of both the dimensionless extrinsic curvature coupling and the background metric. Furthermore in the path integral formalism adopted here where the dynamical variables of the theory are the string's immersion coordinates only, it is shown here explicitly that the volume of the two dimensional diffeomorphism group is cancelled by the volume of the space of tangential fluctuations of the world sheet of the string. This demonstration for onshell amplitudes turns out to be nontrivial.

This paper is organized as follows. In section **0.2**, we consider the classical properties of both the N-G and the extrinsic curvature actions in a curved space-time. We derive their equations of motion and discuss possible solutions. The space-time energy momentum tensor $T^{\mu\nu}$ of the string is derived in section **0.3** and in the appendix we establish that it is covariantly conserved. In section **0.4** we discuss the 1-loop divergences of the theory by integrating over X^μ . As a consequence of reparametrization invariance only the fluctuations normal to the world sheet are dynamical. The longitudinal fluctuations are zero modes and their volume cancels the volume of the diffeomorphism group of the world sheet. It will be seen that a divergent Euler characteristic term appears at the 1 loop level while such a term was previ-

ously known to appear only at the 2 loop level in the conventional approach [2],[25], where the dynamical degrees of freedom are the immersion coordinates and the metric of the world sheet. The condition for 1 loop finiteness is also derived.

0.2 N-G AND EXTRINSIC CURVATURE ACTIONS IN CURVED SPACE-TIME

In the previous section we discussed the N-G action. If the metric $h_{\mu\nu}$ of space-time is taken as dynamical, then, we must include the Hilbert-Einstein action

$$S_{H-E} = -\frac{1}{8\pi G} \int \tilde{R} \sqrt{h} d^D X. \quad (5)$$

We denote the scalar curvature of space-time $\tilde{M}$ by $\tilde{R}$. We use the notation whereby geometric quantities of space-time will be denoted with a tilde and the world sheet quantities will be unadorned.

In order to understand rigid strings, it is necessary to recall the structure equations for immersed surfaces in $\tilde{M}$. First, the equation of Gauss

$$\partial_\alpha \partial_\beta X^\mu + \tilde{\Gamma}_{\nu\rho}^\mu \partial_\alpha X^\nu \partial_\beta X^\rho - \Gamma_{\alpha\beta}^\gamma \partial_\gamma X^\mu = H_{\alpha\beta}^i N^{i\mu}, \quad (6)$$

defines the second fundamental form $H_{\alpha\beta}^i$ ($i = 1, 2, \dots, D-2$). In (6) $\tilde{\Gamma}_{\nu\rho}^\mu$ is the connection in $\tilde{M}$ determined by the metric $h_{\mu\nu}$, while $\Gamma_{\alpha\beta}^\gamma$ is the connection coefficient on string world sheet M determined by the induced metric $g_{\alpha\beta}$ on M . $N^{i\mu}$ ($i = 1, 2, \dots, D-2$) are the $(D-2)$ normals to the string world sheet at $X^\mu(\sigma, \tau)$. We choose the normalization

$$N^{i\mu} N^{j\nu} h_{\mu\nu} = \delta_{ij} \epsilon(i), \quad (7)$$

and

$$\partial_\alpha X^\mu N^{i\nu} h_{\mu\nu} = 0, \quad (i = 1, 2, \dots, D-2). \quad (8)$$

$\epsilon(i) = \pm 1$ depending on whether the world sheet M has an induced metric which is Riemannian or pseudo-Riemannian. For string theory, the world

sheet metric has Lorentzian signature and hence $\epsilon(i)$ in (7) is +1 for $i = 1, 2, \dots, D-2$. In the context of canonical (3+1) gravity, the hypersurface has induced metric which is Riemannian and so $\epsilon(i) = -1$. Note that (6) may be expressed in compact form as

$$\nabla_\alpha \nabla_\beta X^\mu = H_{\alpha\beta}^i N^{i\mu},$$

where ∇_α is the symbol for covariant derivative. Note that $\nabla_\beta X^\mu \equiv \partial_\beta X^\mu$ is a world-vector as well as a 2-vector and hence needs both connections $\tilde{\Gamma}_{\nu\rho}^\mu$ and $\Gamma_{\alpha\beta}^\gamma$ to define its covariant derivative.

Next, we have the equation of Weingarten

$$\partial_\alpha N^{i\mu} + \tilde{\Gamma}_{\nu\rho}^\mu \partial_\alpha X^\nu N^{i\rho} - A_\alpha^{ij} N^{j\mu} = -\epsilon(i) H_\alpha^{i\beta} \partial_\beta X^\mu. \quad (9)$$

In (9)

$$A_\alpha^{ij} = \epsilon(i) \left(N^{j\nu} \tilde{\nabla}_\alpha N^{i\mu} \right) h_{\mu\nu}. \quad (10)$$

$$\tilde{\nabla}_\alpha N^{i\mu} = \partial_\alpha N^{i\mu} + \tilde{\Gamma}_{\nu\rho}^\mu \partial_\alpha X^\nu N^{i\rho}. \quad (11)$$

A_α^{ij} is the 2-d gauge field or gauge connection in the normal bundle.

We shall use the notation whereby we covariantize on all indices. μ is the world index, α, β , the 2-dimensional world sheet index, and i, j the ‘internal’ indices of the normal bundle. Thus

$$\nabla_\alpha N^{i\mu} \equiv \partial_\alpha N^{i\mu} + \tilde{\Gamma}_{\nu\rho}^\mu \partial_\alpha X^\nu N^{i\rho} - A_\alpha^{ij} N^{j\mu}. \quad (12)$$

Note that in D-dimensional space-time $A_\alpha^{ij} = -A_\alpha^{ji}$ are 2-dimensional gauge fields with the gauge group $SO(D-2)$.

In addition to the equations of Gauss and Weingarten, there are three key equations we need in the sequel. These are the equations of Gauss, Codazzi, and Ricci. These are well known and we merely give these below for completeness [26],[27].

Equation of Gauss:

$$\begin{aligned} \tilde{R}_{\mu\nu\rho\sigma} \partial_\alpha X^\mu \partial_\beta X^\nu \partial_\gamma X^\rho \partial_\delta X^\sigma &= R_{\alpha\beta\gamma\delta} \\ &+ \sum_i \epsilon(i) \left(H_{\beta\gamma}^i H_{\alpha\delta}^i - H_{\beta\delta}^i H_{\alpha\gamma}^i \right), \end{aligned} \quad (13)$$

where $R_{\alpha\beta\gamma\delta}$ is the curvature of the string world sheet.

Equation of Codazzi:

$$\nabla_\alpha H_{\beta\gamma}^i - \nabla_\beta H_{\alpha\gamma}^i = \epsilon(i) \tilde{R}_{\mu\nu\rho\sigma} \partial_\alpha X^\mu \partial_\beta X^\nu \partial_\gamma X^\sigma N^{i\rho}, \quad (14)$$

and

Equation of Ricci:

$$\tilde{R}_{\mu\nu\rho\sigma} \partial_\alpha X^\mu \partial_\beta X^\nu N^{j\rho} N^{i\sigma} = F_{\alpha\beta}^{ij} + H_\alpha^{i\gamma} H_{\gamma\beta}^j - H_\beta^{i\gamma} H_{\gamma\alpha}^j, \quad (15)$$

where

$$F_{\alpha\beta}^{ij} = \partial_\alpha A_\beta^{ij} - \partial_\beta A_\alpha^{ij} + A_\alpha^{ik} A_\beta^{kj} - A_\alpha^{jk} A_\beta^{ki}. \quad (16)$$

In (12) we have used the notation

$$\nabla_\alpha H_{\beta\gamma}^i = \partial_\alpha H_{\beta\gamma}^i - \Gamma_{\alpha\beta}^\delta H_{\delta\gamma}^i - \Gamma_{\alpha\gamma}^\delta H_{\beta\delta}^i - A_\alpha^{ij} H_{\beta\gamma}^j. \quad (17)$$

In general these are the only relations between the curvature tensor of the space-time and the extrinsic curvature of the world sheet. In the case when M is a hypersurface in space-time there is only one normal and so $A_\alpha^{ij} \equiv 0$. Further relations may be obtained when space-time metric is written in terms of, say, the isothermal coordinates [11],[28]. The choice of the extrinsic curvature action for strings is motivated from the relation that follows from (13) when $\tilde{R}_{\mu\nu\rho\sigma} = 0$ ($\tilde{M}$ is Minkowski). In this case,

$$R = \sum_i \epsilon(i) \left(H_{\alpha\beta}^i H^{i\alpha\beta} - (H_\alpha^{i\alpha})^2 \right), \quad (18)$$

where R is the scalar curvature of the world sheet. In 2-dimensions

$$\frac{1}{4\pi} \int d\sigma d\tau \sqrt{g} R = \chi, \quad (19)$$

where χ is the Euler characteristic of the two-dimensional manifold. On the other hand each term on the right in (18) is separately reparameterization invariant and hence can be used as the extrinsic curvature action. For convenience, we take this as (subscript R for rigid strings)

$$S_R = \frac{1}{\alpha_0} \int \sqrt{g} H^2 d\sigma d\tau, \quad (20)$$

where

$$H^2 = \sum_i \left(\frac{1}{2} H_\alpha^{i\alpha} \right)^2. \quad (21)$$

We continue to take (20) as the extrinsic curvature action for rigid strings in curved background [From now on we shall drop the factor $\epsilon(i)$ for convenience. In fact for strings all normals will be space-like and $\epsilon(i) = +1$. α_0 in (20) is a dimensionless coupling. In a flat background, it is known that this coupling is asymptotically free [3],[5]. Note that if we considered an action

$$S = \int \tilde{R}_{\mu\nu\rho\sigma} \partial_\alpha X^\mu \partial_\beta X^\nu \partial_\gamma X^\rho \partial_\delta X^\sigma g^{\beta\gamma} g^{\alpha\delta} \sqrt{g} d\sigma d\tau, \quad (22)$$

it will not take into account extrinsic curvature or bending rigidity effects although such terms will probably arise in the effective action starting from the N-G term.

In Minkowski space-time (20), besides being reparameterization invariant on the world sheet, is scale invariant under $X^\mu \rightarrow \lambda X^\mu$. Although not usually emphasized in the literature, the action (20) is actually invariant under space-time conformal transformations [29]-[31]. It is not too difficult to show that (20) is invariant under 4-translations, scale transformation, Lorentz transformations and special conformal transformations [32]

$$X'^\mu = \frac{X^\mu - b^\mu X^2}{1 - 2b \cdot X + b^2 X^2}, \quad (23)$$

where b^μ is a constant D-vector. Invariance under the first three is manifest. It is readily verified that under infinitesimal special conformal transformation, (20) is transformed into a total divergence and hence either vanishes for closed surfaces (without boundary) or turns into a boundary contribution.

Let us concentrate on the properties of (20) in curved background. In order to appreciate the nature of the string world sheet that satisfies the

equation of motion of (20), it is instructive to start with the N-G action (2). A straightforward computation shows that the equation of motion of (2) can be written as

$$\sum_i N^{i\mu} H^i = 0. \quad (24)$$

This is equivalent to $H^i = 0$, $i = 1, 2, \dots, D-2$. In other words surfaces with zero mean curvature, i.e., minimal surfaces, extremize the N-G action. This conclusion is independent of whether the background space-time is flat or curved. Note that the equation of motion for the N-G action is algebraic in the scalar mean curvature. In flat space-time it reduces to

$$(\partial_\tau^2 - \partial_\sigma^2) X^\mu = 0, \quad (25)$$

upon using isothermal coordinates on the string world sheet. In curved background it reads

$$g^{\alpha\beta} \nabla_\alpha \partial_\beta X^\mu = 0, \quad (26)$$

where $\nabla_\alpha \partial_\beta X^\mu$ is defined in (6).

In the path integral formulation, classical solutions provide a starting point for non-perturbative calculations. Thus one can integrate over minimal surfaces for N-G action. In fact the present authors have carried out such a calculation for minimal and harmonic ($H^i = \text{constant}$) surfaces for rigid strings in a flat background space-time and obtained a modified Coulomb gas picture of QCD effective string action [6]. This demonstration [6] however utilized the notion of the Gauss map from the world sheet into the Grassmannian manifold.

Next, consider the equation of motion of rigid string action in curved space-time. Surprisingly, the derivation is rather involved in this case. The special case of an hypersurface in D-dimensions has been discussed in [33]. We sketch below the main steps.

Starting from (20) we have, under $X^\mu \rightarrow X^\mu + \delta X^\mu$,

$$\delta S_R = \frac{1}{\alpha_0} \left(\frac{1}{2} \int \sqrt{g} g^{\alpha\beta} \delta g_{\alpha\beta} H^2 d\sigma d\tau + 2 \int H^i \delta H^i \sqrt{g} d\sigma d\tau \right). \quad (27)$$

A simple calculation yields

$$\begin{aligned}\delta g_{\alpha\beta} = & \nabla_\alpha (\delta X^\mu \partial_\beta X^\nu h_{\mu\nu}) + \nabla_\beta (\delta X^\mu \partial_\alpha X^\nu h_{\mu\nu}) \\ & - 2\delta X^\mu h_{\mu\nu} N^{i\nu} H_{\alpha\beta}^i.\end{aligned}\quad (28)$$

$$\delta N^{i\mu} + \delta X^\sigma N^{i\nu} \Gamma_{\sigma\nu}^\mu = -\partial_\alpha X^\mu \left[\nabla^\alpha (N^{i\nu} \delta X^\sigma h_{\nu\sigma}) + H^{i\alpha\beta} \partial_\beta X^\nu \delta X^\rho h_{\nu\rho} \right]. \quad (29)$$

$$\begin{aligned}\delta H^i = & -\nabla_\alpha \left(H^{i\alpha\beta} \delta X^\mu \partial_\beta X^\nu h_{\mu\nu} \right) + \left(\nabla_\alpha H^{i\alpha\beta} \right) \partial_\beta X^\nu \delta X^\mu h_{\mu\nu} \\ & + H^{i\alpha\beta} H_{\alpha\beta}^j N^{j\nu} \delta X^\mu h_{\mu\nu} + \frac{1}{2} g^{\alpha\beta} \delta H_{\alpha\beta}^i.\end{aligned}\quad (30)$$

$$\begin{aligned}g^{\alpha\beta} H^i \delta H_{\alpha\beta}^i = & \nabla^\alpha \left((\nabla_\alpha \delta X^\mu) N^{i\nu} h_{\mu\nu} H^i - \delta X^\mu h_{\mu\nu} \nabla_\alpha (N^{i\nu} H^i) \right) \\ & + g^{\alpha\beta} H^i \tilde{R}_{\mu\nu\rho\sigma} \partial_\alpha X^\nu \partial_\beta X^\rho \delta X^\sigma N^{i\mu} \\ & + \delta X^\mu h_{\mu\nu} \left[H^i \left(-(\nabla^\alpha H_{\alpha}^{i\gamma}) \partial_\gamma X^\nu - H^i H_{\alpha\beta}^j H^{i\alpha\beta} N^{j\nu} \right) \right. \\ & \left. - 2H^{i\alpha\gamma} (\partial_\gamma X^\nu) \nabla_\alpha H^i + N^{i\nu} \nabla^2 H^i \right].\end{aligned}\quad (31)$$

In (31), $\tilde{R}_{\mu\nu\rho\sigma}$ is the curvature tensor of the background space-time defined by

$$\tilde{R}^\mu{}_{\nu\rho\sigma} = \partial_\rho \tilde{\Gamma}_{\nu\sigma}^\mu - \partial_\sigma \tilde{\Gamma}_{\nu\rho}^\mu + \tilde{\Gamma}_{\nu\sigma}^\lambda \tilde{\Gamma}_{\rho\lambda}^\mu - \tilde{\Gamma}_{\nu\rho}^\lambda \tilde{\Gamma}_{\sigma\lambda}^\mu. \quad (32)$$

Let us note from equation(29) that $\delta N^{i\mu}$ is not a covariant vector, but that $\delta N^{i\mu} + \delta X^\sigma \Gamma_{\sigma\nu}^\mu N^{i\nu}$ is. Using (28)-(31) in (27) one finds

$$\begin{aligned}\delta S_R = & \frac{1}{\alpha_0} \int \{ \nabla^\alpha [(H^2 g_{\alpha\beta} - 2H^i H_{\alpha\beta}^i) \delta X^\mu \partial^\beta X^\nu h_{\mu\nu} \\ & + (\nabla_\alpha \delta X^\mu) N^{i\nu} h_{\mu\nu} H^i - \delta X^\mu h_{\mu\nu} \nabla^\alpha (N^{i\nu} H^i)] \\ & + [(-2(\nabla^\alpha H^i) H^i + H^i \nabla_\beta H^{i\alpha\beta}) \partial_\alpha X^\nu \delta X^\mu h_{\mu\nu} \\ & + (\nabla^2 H^i - 2H^2 H^i + H^j H_{\alpha\beta}^j H^{i\alpha\beta}) \delta X^\mu N^{i\nu} h_{\mu\nu} \\ & - g^{\alpha\beta} H^i \partial_\alpha X^\nu \partial_\beta X^\rho N^{i\mu} \delta X^\sigma \tilde{R}_{\mu\nu\rho\sigma}] \} \sqrt{g} d\sigma d\tau.\end{aligned}\quad (33)$$

Equations of motion can be obtained by decomposing δX^μ into normal and tangential variations of the world sheet. Accordingly, we write

$$\delta X^\mu = \xi_j N^{j\mu} + \partial_\alpha X^\mu \xi^\alpha. \quad (34)$$

Then it follows from (33) that the equation of motion for normal variations is

$$\nabla^2 H^i - 2H^2 H^i + H^j H^{j\alpha\beta} H_{\alpha\beta}^i - g^{\alpha\beta} H^j \tilde{R}_{\mu\nu\rho\sigma} \partial_\alpha X^\nu \partial_\beta X^\rho N^{j\mu} N^{i\sigma} = 0, \quad (35)$$

where $i = 1, 2, \dots, D-2$. The tangential variations yield

$$\sum_i H^i [\nabla_\beta H^{i\alpha\beta} - 2\nabla^\alpha H^i - \tilde{R}_{\mu\nu\rho\sigma} \partial_\gamma X^\nu \partial_\delta X^\rho g^{\gamma\delta} \partial^\alpha X^\sigma N^{i\mu}] = 0. \quad (36)$$

The equations of motion for the action that is a sum of the N-G and extrinsic curvature actions is easily seen to be

$$\begin{aligned} \nabla^2 H^i - 2H^i(H^2 + k) + H^j H^{j\alpha\beta} H_{\alpha\beta}^i \\ - g^{\alpha\beta} H^j \tilde{R}_{\mu\nu\rho\sigma} \partial_\alpha X^\nu \partial_\beta X^\rho N^{j\mu} N^{i\sigma} = 0, \end{aligned} \quad (37)$$

where $k = -\frac{\alpha_0}{2\pi\alpha'}$. Equation (36) is however unchanged. In (35)

$$\begin{aligned} \nabla^2 &\equiv g^{\alpha\beta} \nabla_\alpha \nabla_\beta; \\ \nabla_\alpha H^i &= \partial_\alpha H^i - A_\alpha^{ij} H^j. \end{aligned} \quad (38)$$

Equation (36) is actually not an equation of motion. It is just the structure equation of Codazzi (14) once contracted. This is just a consequence of reparametrization invariance of the action. Thus the surfaces with mean curvature H^i satisfying (35) extremize the rigid string action in curved background. It is worth remarking here that the extrinsic curvature action contains fourth derivative operator acting on X^μ . Nevertheless its equation of motion is expressible as a second order non linear partial differential equation in the mean scalar curvature H^i of the world sheet. Once one has a

solution of equation (35) for H^i then one can solve for immersion coordinates X^μ from equation (6). Solution of equation of motion for N-G strings in cosmological space-times [1] reveals stable, dual to stable and unstable solutions. In the language employed here, minimal surfaces are still solutions to N-G string in curved space-times. Hence the result of [1] indicates that not all minimal surfaces are stable immersions in space-time. It would thus be interesting to investigate similar stability considerations for rigid strings. In flat space-time, solutions to (35) in 3-dimensional Euclidean space are referred to as Wilmore surfaces [31] and have been investigated in the context of biological membranes in [34]. This equation takes the form (after using (13))

$$\nabla^2 H + 2H(H^2 - \frac{R}{2}) = 0, \quad (39)$$

where R is the scalar curvature of the world sheet. A sphere satisfies this equation with constant H . Solutions with non-zero genus are known to exist for (39) [31]. Note that (39) is a nonlinear wave equation for H^i in the context of string theory.

The special case of a hypersurface in curved space-time is worth considering. In this case the indices i, j take only one value and α, β take values 1, 2, 3. Then (35) reads

$$\nabla^2 H - 2H^3 + HH_{\alpha\beta}H^{\alpha\beta} - H\tilde{R}_{\mu\nu\rho\sigma}\partial_\alpha X^\nu\partial_\beta X^\rho g^{\alpha\beta}N^\mu N^\sigma = 0. \quad (40)$$

Using (13) to eliminate $H_{\alpha\beta}H^{\alpha\beta}$ from (40) we find

$$\begin{aligned} \nabla^2 H + 2H^3 + H[\tilde{R}_{\mu\nu\rho\sigma}\partial_\alpha X^\mu\partial_\beta X^\nu\partial_\gamma X^\rho\partial_\delta X^\sigma g^{\alpha\gamma}g^{\beta\delta} \\ - R - \tilde{R}_{\mu\nu\rho\sigma}\partial_\alpha X^\nu\partial_\beta X^\rho g^{\alpha\beta}N^\mu N^\sigma] = 0. \end{aligned} \quad (41)$$

In [33] the term $\tilde{R}_{\mu\nu\rho\sigma}\partial_\alpha X^\mu\partial_\beta X^\nu\partial_\gamma X^\rho\partial_\delta X^\sigma g^{\alpha\gamma}g^{\beta\delta}$ is replaced by $\tilde{R}$, which in their notation presumably is the scalar curvature of $\tilde{M}$, which is given by $\tilde{R}_{\mu\nu\rho\sigma}h^{\mu\rho}h^{\nu\sigma}$. This would imply that $\partial_\alpha X^\mu\partial_\gamma X^\rho g^{\alpha\gamma} = h^{\mu\rho}$ which is, of course, wrong [see equation (57) below].

In string theory, it would be most interesting to investigate solutions to (35) in cosmological backgrounds. Let us however restrict here to de Sitter

or anti-de Sitter space-times, i.e., space-times of constant curvature. We can write

$$\tilde{R}_{\mu\nu\rho\sigma} = K(h_{\mu\rho}h_{\nu\sigma} - h_{\mu\sigma}h_{\nu\rho}). \quad (42)$$

The equation of motion (35) reads (we choose $\epsilon(i) = 1$)

$$\nabla^2 H^i - 2H^i(H^2 + K) + H^j H_{\alpha\beta}^j H^{i\alpha\beta} = 0. \quad (43)$$

It would be of interest to analyze solutions to (43).

Locally conserved currents arising as a consequence of the invariance of the extrinsic curvature action under local space-time conformal transformations may be readily obtained from the total divergence terms in (33). Let us discuss, in particular only the string's space-time momenta. It is easily seen to be given by

$$(P_\mu^\alpha)_R = \frac{\sqrt{g}}{\alpha_0} [(H^2 g^{\alpha\beta} - H^i H^{i\alpha\beta}) \partial_\beta X^\nu h_{\mu\nu} - (\nabla^\alpha H^i) N^{i\nu} h_{\mu\nu}]. \quad (44)$$

We notice that $(P_\mu^\alpha)_R N^{i\mu} = -\frac{\sqrt{g}}{\alpha_0} (\nabla^\alpha H^i)$ is the component of string momenta normal to its world sheet. This measures the bending energy of the rigid string. In comparison the space-time momentum of the N-G string is given by

$$(P_\mu^\alpha)_{N-G} = \frac{\sqrt{g}}{2\pi\alpha'} g^{\alpha\beta} (\partial_\beta X^\nu) h_{\mu\nu} \quad (45)$$

which has no component normal to the world sheet.

0.3 STRING ENERGY-MOMENTUM TENSOR

The string's space-time energy-momentum tensor $T^{\mu\nu}$ is obtained by taking the functional derivative of the action with respect to the metric $h_{\mu\nu}$ at the space-time point X. The calculation nevertheless turns out to be rather long. We quote here the result.

$$\delta S_R = \frac{1}{2} \int \sqrt{h} T^{\mu\nu} \delta h_{\mu\nu} d^D X, \quad (46)$$

with

$$\begin{aligned}
T^{\mu\nu}(X) = & \frac{1}{\alpha_0} \int d\sigma d\tau \frac{\delta^D(X - X(\sigma, \tau))}{\sqrt{h}} \{ \sqrt{g} \partial_\alpha X^\mu \partial_\beta X^\nu \\
& (H^2 g^{\alpha\beta} - H^i H^{i\alpha\beta}) \\
& - \frac{1}{2} \sqrt{g} g^{\alpha\beta} \nabla_\beta H^i (N^{i\nu} \partial_\alpha X^\mu + N^{i\mu} \partial_\alpha X^\nu) \} \\
& - \frac{\nabla_\rho}{2\alpha_0} \int d\sigma d\tau \frac{\delta^D(X - X(\sigma, \tau))}{\sqrt{h}} \sqrt{g} g^{\alpha\beta} H^i [\partial_\beta X^\mu (N^{i\nu} \partial_\alpha X^\rho \\
& - N^{i\rho} \partial_\alpha X^\nu) + \partial_\beta X^\nu (N^{i\mu} \partial_\alpha X^\rho - N^{i\rho} \partial_\alpha X^\mu)]. \quad (47)
\end{aligned}$$

Note that X in (47) is just a space-time point, whereas $X(\sigma, \tau)$ stands for the string dynamical variables. One sees from the D-dimensional Dirac delta function in (47) that $T^{\mu\nu}(X)$ vanishes unless X is exactly on the string world-sheet. It is instructive to compare (47) with the energy-momentum tensor of the N-G strings.

$$T_{N-G}^{\mu\nu}(X) = \frac{1}{\alpha'} \int d\sigma d\tau \frac{\delta^D(X - X(\sigma, \tau))}{\sqrt{h}} \sqrt{g} g^{\alpha\beta} \partial_\alpha X^\mu \partial_\beta X^\nu. \quad (48)$$

Note that the flux of $T_{N-G}^{\mu\nu}$ along the normal direction $T_{N-G}^{\mu\nu} N_\nu^i(X) = 0$. This is no longer true for rigid strings as can be seen from (47). This is consistent with the remarks in the introduction that bending of the strings costs energy.

It can be verified, although the calculations are rather long, that

$$\nabla_\mu T^{\mu\nu} = 0 \quad (49)$$

when the equation of motion are taken into account. Demonstration of (49) is sketched in the Appendix.

0.4 1-LOOP QUANTUM EFFECTS

We now discuss 1-loop quantum effects for strings in background space-time. In the conventional treatment, the N-G action is taken in the form (1) in which the dynamical variables are $g_{\alpha\beta}$ and X^μ and in this form the action

is regarded as a non linear σ model [2]. We show in this section that it is more convenient to perform computations by regarding the world sheet as an immersed surface and the metric on it as induced from the background space-time. The dynamical fields are just the immersion coordinates X^μ . It will become clear that in this scheme not only the metric of the world sheet but also its extrinsic geometry play a key role in the dynamics. The partition function for the string is given by

$$\mathbf{Z} = \int \frac{\mathcal{D}X^\mu}{Vol} e^{-\frac{1}{2\pi\alpha'} \int \sqrt{g} d\sigma d\tau}, \quad (50)$$

where Vol is the volume of the 2-d diffeomorphism group. To calculate 1-loop effects we need the second variation of the action. We now evaluate this on-shell, i.e. for minimal surfaces. As in (34) we decompose the fluctuations δX^μ into normal and tangential components. For the N-G action the first variation of the action can be written as

$$\delta S = -\frac{1}{\pi\alpha'} \int \sqrt{g} H^i \xi_i d\sigma d\tau. \quad (51)$$

To evaluate the second variation on-shell ($H^i = 0$), we need only evaluate the variation of H^i . It is convenient to rewrite (30) and (31) in the following form

$$\begin{aligned} \delta H_{\alpha\beta}^i &= (\nabla_\alpha \nabla_\beta \delta_j^i - H_{\alpha\gamma}^i H_\beta^{i\gamma} \\ &\quad - \tilde{R}_{\mu\nu\rho\sigma} N^{i\mu} N^{j\sigma} \partial_\alpha X^\rho \partial_\beta X^\nu) \xi_j \\ &\quad + (H_\alpha^{i\gamma} \nabla_\beta + H_\beta^{i\gamma} \nabla_\alpha) \xi_\gamma + (\nabla^\gamma H_{\alpha\beta}^i) \xi_\gamma. \end{aligned} \quad (52)$$

In arriving at the result above from (30) and (31), we have made use of the equation of Gauss (13). From above we find that δH^i can be expressed as follows,

$$\delta H^i = \frac{1}{2} \mathbf{O}_j^i \xi^j + (\nabla^\alpha H^i) \xi_\alpha, \quad (53)$$

where

$$\mathbf{O}_j^i = \nabla^2 \delta_j^i + H_{\alpha\beta}^i H^{j\alpha\beta} - \tilde{R}_{\mu\nu\rho\sigma} g^{\alpha\beta} \partial_\alpha X^\nu \partial_\beta X^\rho N^{i\mu} N^{j\sigma}. \quad (54)$$

Substituting from above we can write the second variation of (2) as follows

$$\delta^2 S = -\frac{1}{2\pi\alpha'} \int \sqrt{g} \xi^i \mathbf{O}_j^i \xi^j d\sigma d\tau. \quad (55)$$

Note that only the normal-normal fluctuations appear in the second variation when evaluated on-shell. As a consequence $\int \frac{\mathcal{D}X^\mu}{Vol} \rightarrow \int \frac{\mathcal{D}\xi^\alpha}{Vol} \mathcal{D}\xi^i \rightarrow \int \mathcal{D}\xi^i$.

The logarithmic correction to the effective action can now be read off from (54) and (55) [35]

$$\begin{aligned} \Gamma^{(1)} = & \frac{I}{2} \int d\sigma d\tau \sqrt{g} \{ \tilde{R}_{\lambda\nu\rho\sigma} g^{\alpha\beta} \partial_\alpha X^\nu \partial_\beta X^\rho N^{i\lambda} N^{i\sigma} \\ & - H_{\alpha\beta}^i H^{i\alpha\beta} \}, \end{aligned} \quad (56)$$

where

$$I = \int^\Lambda \frac{d^2 k}{(2\pi)^2 k^2}. \quad (57)$$

The covariant derivative ∇ in (54) contains the gauge connection A_α^{ij} . But counter terms dependent on A_α^{ij} do not arise since they can appear only through $Tr(F_{\alpha\beta} F^{\alpha\beta})$ which is a dimension four operator.

Now from Gauss' formula (13) we find

$$H_{\alpha\beta}^i H^{i\alpha\beta} = 4H^2 + R - \tilde{R}_{\mu\nu\rho\sigma} \partial_\alpha X^\mu \partial_\beta X^\nu \partial_\gamma X^\rho \partial_\delta X^\sigma g^{\beta\gamma} g^{\alpha\delta}. \quad (58)$$

R in above is the scalar curvature of the world sheet. Note that for N-G action we put $H = 0$ in the above formula. Substituting (58) in (56) and making use of the completeness relation

$$\sum_{i=1}^{D-2} N^{i\lambda} N^{i\sigma} + \partial_\gamma X^\lambda \partial_\delta X^\sigma g^{\gamma\delta} = h^{\lambda\sigma}, \quad (59)$$

we get

$$\Gamma^{(1)} = \frac{I}{2} \int d\sigma d\tau \sqrt{g} \{ \tilde{R}_{\nu\rho} \partial_\alpha X^\nu \partial_\beta X^\rho g^{\alpha\beta} - R \} \quad (60)$$

where $\tilde{R}_{\nu\rho}$ is the Ricci tensor of space-time. As claimed before, the Euler characteristic term appears in 1-loop order. This is to be contrasted with the results in [2], [25]. Both divergent terms in (60) can be absorbed in the renormalization of the background metric $h_{\mu\nu}$. Thus, defining renormalized $h_{\mu\nu}$ by

$$h_{\mu\nu}^R = h_{\mu\nu} + \delta h_{\mu\nu}, \quad (61)$$

we find that

$$\begin{aligned} \sqrt{g} &= \det(\partial_\alpha X^\mu \partial_\beta X^\nu (h_{\mu\nu}^R - \delta h_{\mu\nu})) \\ &\simeq \sqrt{g} (1 - \frac{1}{2} g^{\alpha\beta} \partial_\alpha X^\mu \partial_\beta X^\nu \delta h_{\mu\nu}). \end{aligned} \quad (62)$$

Thus, we define

$$\delta h_{\mu\nu} = 2\pi\alpha' I [\tilde{R}_{\mu\nu} - \frac{1}{2} R h_{\mu\nu}]. \quad (63)$$

The divergent Euler characteristic term thus gets absorbed in the redefinition of the background metric rather than in Liouville field as happens in the conventional procedure [2]. Thus, for 1-loop finiteness of string theory the necessary condition is

$$\tilde{R}_{\mu\nu} - \frac{1}{2} R h_{\mu\nu} = 0 \quad (64)$$

which is different from the Ricci flatness condition met in the sigma models. Note, also, that R in (64) is the scalar curvature of the 2-d world sheet of the string. When $R = 0$, (64) is the vacuum Einstein equation and this agrees with the results obtained in [2], [4] using the formalism where both the 2-d metric and immersion coordinates are treated dynamically. There this equation is derived for flat world sheet. The crucial difference from their results arises from the scalar curvature of the world sheet appearing in (64). In [2], it is shown that the divergent Euler characteristic term appears in the second order perturbation and it is absorbed in the renormalization of the Liouville mode. We note here that, in the formalism used here, this term appears in the first order and is absorbed in the background metric renormalization. Eqn.(64) has a very nice interpretation. First note that as

a consequence of (64) $\tilde{R} = (D/2)R$. This makes sense for, when $D=2$, $\tilde{R}$ evaluated on the world sheet must coincide with the scalar curvature of the world sheet. Using this relation back in (64) we can rewrite it as

$$\tilde{R}_{\mu\nu} - \frac{1}{2}\tilde{R}h_{\mu\nu} = \frac{2-D}{4}Rh_{\mu\nu},$$

which can be interpreted as Einstein equation with a varying cosmological constant $\Lambda = \frac{2-D}{4}R$.

We now calculate 1-loop effects for the extrinsic curvature action. The coupling constant α_0 in (20) is asymptotically free in flat space-time. So, in this case we will encounter renormalization of both the space-time metric and the coupling of the extrinsic curvature action. As in the case of the N-G action, we start by calculating the second variation of the extrinsic geometric action. This, however turns out to be rather involved and we sketch the essential steps below. The first variation of this action given in (33) is, after ignoring the total divergence terms and the longitudinal fluctuations terms, (which vanish due to the equation of Gauss) can be written as

$$\begin{aligned} \delta S_R = & \int \sqrt{g}(\nabla^2 H^i - 2H^2 H^i + H^j H_{\alpha\beta}^j H^{i\alpha\beta} \\ & - H^j \tilde{R}_{\lambda\nu\rho\sigma} \partial_\alpha X^\nu \partial_\beta X^\rho g^{\alpha\beta} N^{j\lambda} N^{i\sigma}) \xi_i(\sigma, \tau) d\sigma d\tau, \end{aligned} \quad (65)$$

where $\xi_i(\sigma, \tau)$ is the normal fluctuation defined in (34). Starting from the above equation we evaluate its on-shell variation. Thus, it is sufficient to evaluate the variation of the terms within the brackets. This is because of the equation of motion (35). It is clear that there will be terms that involve operators that couple only normal fluctuations and a piece that couples tangential fluctuations to the normal fluctuations. It will turn out that these second type of terms vanish as a consequence of the equation of motion (35). The calculations needed to evaluate the second variation of the extrinsic curvature action are lengthy. We give below the key steps.

In addition to the results used in the second variation of the N-G action, as well as equations (27), (28), (52), and (53), we need the following results: The variation of $\Gamma_{\alpha\beta}^\gamma$:

$$\delta \Gamma_{\alpha\beta}^\gamma = \frac{1}{2}g^{\gamma\delta}(\nabla_\alpha \nabla_\beta + \nabla_\beta \nabla_\alpha)\xi_\delta - \frac{1}{2}g^{\gamma\delta}R^\epsilon{}_{\beta\alpha\delta}\xi_\epsilon$$

$$\begin{aligned}
& -\frac{1}{2}g^{\gamma\delta}R^\epsilon{}_{\alpha\beta\delta}\xi_\epsilon - g^{\gamma\delta}\nabla_\alpha(H^j_{\delta\beta}\xi^j) \\
& -g^{\gamma\delta}\nabla_\beta(H^j_{\alpha\delta}\xi^j) + g^{\gamma\delta}\nabla_\delta(H^j_{\alpha\beta}\xi^j).
\end{aligned} \tag{66}$$

In (66), $R^\epsilon{}_{\beta\alpha\delta}$ is the curvature tensor of the world sheet. The longitudinal fluctuations are denoted by ξ with greek indices while the normal fluctuations are denoted by ξ with latin indices. Next, the variation of the normal gauge connection A^{ij}_α : From its definition

$$A^{ij}_\alpha = N^{j\nu}(\tilde{\nabla}_\alpha N^{i\mu})h_{\mu\nu}, \tag{67}$$

where $\tilde{\nabla}_\alpha$ is defined in (11), it can be verified that

$$\begin{aligned}
\delta A^{ij}_\alpha &= -F^{ij}_{\alpha\beta}\xi^\beta + (H^i_{\alpha\beta}\nabla^\beta\xi^j - H^j_{\alpha\beta}\nabla^\beta\xi^i) \\
&+ \tilde{R}_{\mu\sigma\lambda\rho}N^{i\sigma}N^{j\mu}\partial_\alpha X^\rho N^{k\lambda}\xi^k.
\end{aligned} \tag{68}$$

In (86), $F^{ij}_{\alpha\beta}$ is the field strength of A^{ij}_α defined in (16).

We now give the results for the variation of each of the terms in (65).

$$\begin{aligned}
\delta(\nabla^2 H^i) &= (\nabla_\gamma(\nabla^2 H^i))\xi^\gamma + \frac{1}{2}\nabla^2(\mathbf{O}^i_j\xi^j) + 2(\nabla^\alpha\nabla^\beta H^i)H^j_{\alpha\beta}\xi^j \\
&- 2(\nabla^\alpha H^j)[(H^i_{\alpha\beta}\nabla^\beta\xi^j - H^j_{\alpha\beta}\nabla^\beta\xi^i) \\
&+ \tilde{R}_{\mu\sigma\lambda\rho}N^{i\sigma}N^{j\mu}\partial_\alpha X^\rho N^{k\lambda}\xi^k] \\
&- H^j\nabla^\alpha[\tilde{R}_{\mu\sigma\lambda\rho}N^{i\sigma}N^{j\mu}\partial_\alpha X^\rho N^{k\lambda}\xi^k \\
&+ (H^i_{\alpha\beta}\nabla^\beta\xi^j - H^j_{\alpha\beta}\nabla^\beta\xi^i)] \\
&+ 2(\nabla^\gamma H^i)[\nabla^\alpha(H^j_{\alpha\gamma}\xi^j) - \nabla_\gamma(H^j\xi^j)].
\end{aligned} \tag{69}$$

The variation of $(H^2 H^i)$ is given by

$$\delta(H^2 H^i) = (\nabla^\alpha(H^2 H^i))\xi_\alpha + (\frac{1}{2}H^2\delta^i_j + H^i H^j)\mathbf{O}^i_j\xi^k, \tag{70}$$

and

$$\begin{aligned}
\delta(H^j H^j_{\alpha\beta} H^{i\alpha\beta}) &= (\nabla^\alpha(H^j H^j_{\beta\gamma} H^{i\beta\gamma}))\xi_\alpha + \frac{1}{2}H^j_{\alpha\beta} H^{i\alpha\beta}\mathbf{O}^j_k\xi^k \\
&+ H^j H^{i\alpha\beta}\mathbf{O}^j_{k,\alpha\beta}\xi^k + H^j H^{j\alpha\beta}\mathbf{O}^i_{k,\alpha\beta}\xi^k,
\end{aligned} \tag{71}$$

where

$$\mathbf{O}_{k,\alpha\beta}^i = \nabla_\alpha \nabla_\beta \delta_k^i + H_{\alpha\gamma}^i H_{k\beta}^\gamma - \tilde{R}_{\mu\nu\rho\sigma} N^{i\mu} N_k^\sigma \partial_\alpha X^\rho \partial_\beta X^\nu. \quad (72)$$

The above operator $\mathbf{O}_{k,\alpha\beta}^i$ satisfies

$$\mathbf{O}_{k,\alpha\beta}^i g^{\alpha\beta} = \mathbf{O}_k^i,$$

which is defined in (54).

The final variation we need is given below.

$$\begin{aligned} \delta(g^{\alpha\beta} H^j \tilde{R}_{\mu\nu\rho\sigma} \partial_\alpha X^\nu \partial_\beta X^\rho N^{j\mu} N^{i\sigma}) &= \nabla^\alpha (H^j \tilde{R}_{\lambda\nu\rho\sigma} \partial_\alpha X^\nu \partial_\beta X^\rho g^{\alpha\beta} N^{j\lambda} \\ &\quad N^{i\sigma}) \xi_\alpha \\ &\quad + H^j (\nabla_\mu \tilde{R}_{\lambda\nu\rho\sigma}) \partial_\alpha X^\nu \partial_\beta X^\rho g^{\alpha\beta} N^{j\lambda} \\ &\quad N^{i\sigma} N^{k\mu} \xi^k \\ &\quad + \frac{1}{2} \tilde{R}_{\lambda\nu\rho\sigma} \partial_\alpha X^\nu \partial_\beta X^\rho g^{\alpha\beta} N^{j\lambda} N^{i\sigma} \mathbf{O}_k^i \xi^k \\ &\quad + H^j \tilde{R}_{\lambda\nu\rho\sigma} N^{j\lambda} N^{i\sigma} (\partial_\alpha X^\rho N^{k\nu} \\ &\quad + \partial_\alpha X^\nu N^{k\rho}) \nabla^\alpha \xi^k \\ &\quad + 2H^j (\nabla_\gamma H^i) \nabla^\gamma \xi^j + 2H^j (\nabla_\gamma H^j) \\ &\quad \nabla^\gamma \xi^i \\ &\quad - H^j (\nabla^\alpha H_{\alpha\gamma}^i) \nabla^\gamma \xi^j - H^j (\nabla^\alpha H_{\alpha\gamma}^j) \\ &\quad \nabla^\gamma \xi^i. \end{aligned} \quad (73)$$

Making use of the above results we can write down the second variation of the extrinsic curvature action in background space-time. Notice that the terms that couple the normal fluctuations to the tangential ones vanish because of the equation of motion (35). We thus find for $\delta^2 S$ the following:

$$\delta^2 S = \frac{1}{\alpha_0} \int (\xi^i \tilde{\mathbf{O}}_k^i \xi^k) \sqrt{g} d\sigma d\tau, \quad (74)$$

where

$$\begin{aligned} \tilde{\mathbf{O}}_k^i &= \frac{1}{2} \mathbf{O}_j^i \mathbf{O}_k^j - (H^2 \delta_j^i + 2H^i H^j) \mathbf{O}_k^j + H^j H^{i\alpha\beta} \mathbf{O}_{k,\alpha\beta}^j \\ &\quad + H^j H^{j\alpha\beta} \mathbf{O}_{k,\alpha\beta}^i + 2(\nabla^\alpha \nabla^\beta H^i) H_{\alpha\beta}^k \end{aligned}$$

$$\begin{aligned}
& -2\nabla^\alpha H^j (H_{\alpha\beta}^i \nabla^\beta \delta_k^j - H_{\alpha\beta}^j \nabla^\beta \delta_k^i) \\
& -2(\nabla^\alpha H^j) \tilde{R}_{\mu\sigma\lambda\rho} N^{i\sigma} N^{j\mu} \partial_\alpha X^\rho N^{k\lambda} \\
& -H^j (\nabla^\alpha [H_{\alpha\beta}^i \nabla^\beta \delta_k^j - H_{\alpha\beta}^j \nabla^\beta \delta_k^i + \tilde{R}_{\mu\sigma\lambda\rho} N^{i\sigma} N^{j\mu} \partial_\alpha X^\rho N^{k\lambda}]) \\
& +2\nabla^\gamma H^i [\nabla^\alpha H_{\alpha\gamma}^j + H_{\alpha\gamma}^j \nabla^\alpha - \nabla^\gamma H^j - H^j \nabla^\gamma] \delta_k^j \\
& -H^j (\nabla_\mu \tilde{R}_{\lambda\nu\rho\sigma}) \partial_\alpha X^\nu \partial_\beta X^\rho g^{\alpha\beta} N^{j\lambda} N^{k\mu} - 2H^j (\nabla_\gamma H^i) \nabla^\gamma \delta_k^j \\
& -2H^j (\nabla_\gamma H^j) \nabla^\gamma \delta_k^i + H^j (\nabla^\alpha H_{\alpha\gamma}^i) \nabla^\gamma \delta_k^j + H^j (\nabla^\alpha H_{\alpha\gamma}^j) \nabla^\gamma \delta_k^i. \quad (75)
\end{aligned}$$

It is straightforward to add the second variation of the N-G action to the above. Again, because of the equation of motion for the combined action given in (37) the second variation is simply the sum of the two variations.

One loop divergent terms can be readily read off from (75). The bare propagator for the normal fluctuations is

$$\langle \xi^i \xi^j \rangle = \delta_j^i \frac{1}{p^4 + (2\pi\alpha')^{-1} p^2} \quad (76)$$

As observed in the case of the N-G action, divergent terms involving A_α^{ij} do not arise since they can only appear through $Tr(F^{\alpha\beta} F_{\alpha\beta})$ which is a dimension four operator. There are two types of divergent terms which are readily computed to give the following result.

$$\begin{aligned}
\Gamma^1 &= -DI \int H^2 \sqrt{g} d\sigma d\tau \\
& -I \int \left(\frac{1}{2} R h_{\nu\rho} - \tilde{R}_{\nu\rho} \right) \partial_\alpha X^\nu \partial_\beta X^\rho g^{\alpha\beta} \sqrt{g} d\sigma d\tau, \quad (77)
\end{aligned}$$

where I is as given in (57). Divergent extrinsic curvature terms arise from terms 1 to 4 and 8 in (75), whose sum yields the factor D in (77). In writing Γ^1 in the form (77), we have made use of the equation of Gauss given in (58).

It is seen that the first term in (77) can be absorbed in the redefinition of the extrinsic curvature coupling α_0 according as

$$\alpha_R = \frac{\alpha_0}{1 - \alpha_0 DI}. \quad (78)$$

Notice that the factor D appears in the denominator in (78), which is the dimensionality of the background space-time, rather than $D - 2$ that would

be expected of a non linear sigma model in which the sigma model fields are the $D - 2$ normals to the surface. We have thus verified that the coupling for the extrinsic curvature action in an arbitrary curved background is asymptotically free. We have avoided using the sigma model formalism and worked directly in terms of the immersion coordinates X^μ as the only dynamical degrees of the theory. The other 1-loop divergent term in (77) can be absorbed, as in the case of the N-G action, by renormalizing the background metric exactly as before. We have not calculated the finite contributions to the partition function in this paper. The simplest case would be to integrate over minimal surfaces. We can take minimal surfaces with punctures which can be interpreted as locations of instanton quarks [6].

0.5 SUMMARY

In this paper we have explored both the intrinsic and extrinsic geometric properties of strings immersed in a D dimensional curved space-time. We have used the formalism in which only the string coordinates X^μ are the dynamical fields, while the metric on the world sheet is induced from the string's embedding in space-time. This procedure allows one to take full advantage of the equations of Gauss, Codazzi, and Ricci for immersed surfaces in a general background space-time. It is also established that the extrinsic geometry of the strings plays as important a role as the Nambu-Goto term in determining the dynamics of the strings. We have studied the classical equations of motion and the energy-momentum tensor of strings described by both the N-G and extrinsic curvature actions. The equations of motion determine the scalar mean curvature of the immersed world sheets. We have derived an expression for the covariantly conserved energy-momentum tensor for rigid strings. We have also studied 1-loop quantum corrections using the background field method. The second variation of the action when calculated on-shell couples only the normal fluctuations of the world sheet. The tangential fluctuations (zero modes) decouple from the normal fluctuation modes. We have discussed the renormalization of the coupling constant of the extrinsic curvature action as well as the renormalization of the background metric. 1-loop finiteness requires a vacuum Einstein type equation.

Acknowledgements:

We thank R.Capovilla and J.Guven for their comments on the original version of the manuscript. This work is supported in part by an operating grant from the Natural Sciences and Engineering Research Council of Canada.

APPENDIX

From $T^{\mu\nu}$ defined in (47) we demonstrate below that $\nabla_\mu T^{\mu\nu} = 0$, when the equations of motion (35) are satisfied. A direct calculation shows that

$$\begin{aligned}
\nabla_\mu T^{\mu\nu} = & \int d\sigma d\tau \frac{\delta^D(X - X(\sigma, \tau))}{\sqrt{h}} \sqrt{g} \{ [-\nabla^2 H^i + 2H^2 H^i - H^j H_{\alpha\beta}^i H^{j\alpha\beta}] \\
& N^{i\nu} + H^i \partial_\beta X^\nu [2g^{\alpha\beta} \nabla_\alpha H^i - \nabla_\alpha H^{i\alpha\beta}] \\
& + \frac{1}{2} ((\nabla^2 H^i) N^{i\nu} - (\nabla_\alpha H^i) H^{i\alpha\beta} \partial_\beta X^\nu) \} \\
& - \frac{1}{2} \nabla_\mu \int d\sigma d\tau \frac{\delta^D(X - X(\sigma, \tau))}{\sqrt{h}} \sqrt{g} g^{\alpha\beta} (\nabla_\beta H^i) N^{i\mu} \partial_\alpha X^\nu \\
& - [\nabla_\mu, \nabla_\rho] \int \frac{\delta^D(X - X(\sigma, \tau))}{\sqrt{h}} \sqrt{g} g^{\alpha\beta} H^i \partial_\beta X^\nu \partial_\alpha X^\rho N^{i\mu} \\
& - \frac{1}{2} \nabla_\rho \nabla_\mu \int \frac{\delta^D(X - X(\sigma, \tau))}{\sqrt{h}} \sqrt{g} g^{\alpha\beta} H^i \partial_\beta X^\mu \\
& (N^{i\nu} \partial_\alpha X^\rho - N^{i\rho} \partial_\alpha X^\nu). \tag{A-1}
\end{aligned}$$

The last term in (A-1) cancels the second and third terms. Finally considering the 2-dimensional integral in the commutator term as a tensor $A^{\nu\rho\mu}(X)$ and applying the rule

$$[\nabla_\mu, \nabla_\rho] A^{\nu\rho\mu} = R_{\sigma\mu\rho}^\nu A^{\sigma\rho\mu} + R_{\sigma\mu\rho}^\rho A^{\nu\sigma\mu} + R_{\sigma\mu\rho}^\mu A^{\nu\rho\sigma} \tag{A-2}.$$

and taking advantage of the Dirac delta function under the integral, we can write $\nabla_\mu T^{\mu\nu}$ as

$$\begin{aligned}
\nabla_\mu T^{\mu\nu} = & \int d\sigma d\tau \frac{\delta^D(X - X(\sigma, \tau))}{\sqrt{h}} \sqrt{g} \{ [-\nabla^2 H^i + 2H^2 H^i \\
& - H^j H_{\alpha\beta}^i H^{j\alpha\beta}] N^{i\nu} + H^i [2\nabla^\beta H^i - \nabla_\alpha H^{i\alpha\beta}] \partial_\beta X^\nu \\
& + H^i \tilde{R}^\nu{}_{\sigma\rho\mu} \partial_\alpha X^\sigma \partial_\beta X^\rho g^{\alpha\beta} N^{i\mu} \}. \tag{A-3}
\end{aligned}$$

Requiring $\nabla_\mu T^{\mu\nu} = 0$, we find that

$$\begin{aligned}
& (\nabla^2 H^i - 2H^2 H^i + H^j H_{\alpha\beta}^i H^{j\alpha\beta}) N^{i\nu} \\
& - H^i \tilde{R}^\nu{}_{\sigma\rho\mu} \partial_\alpha X^\sigma \partial_\beta X^\rho g^{\alpha\beta} N^{i\mu} - H^i [2\nabla^\beta H^i - \nabla_\alpha H^{i\alpha\beta}] \partial_\beta X^\nu = 0. \tag{A-4}
\end{aligned}$$

Multiplying (A-4) by $N^{k\mu} h_{\mu\nu}$ we find this condition is the equation of motion (35) while the tangential projection yields the Codazzi equation (36).

References

- [1] H.J.de Vega and N.Sanchez, preprint- hep-th/9512074 (1995)
- [2] A.M.Polyakov in Gauge Fields and Strings, Harwood Academic publishers (1987).
- [3] A.M.Polyakov, Nucl.Phys. **B268**, 406 (1986).
- [4] M.B.Green, J.H.Schwarz and E.Witten, STRING THEORY, Vol.I and II. Cambridge University Press, Cambridge (1987).
- [5] K.S.Viswanathan, R.Parthasarathy, and D.Kay Ann.Phys. **206**, 237 (1991).
- [6] K.S.Viswanathan and R.Parthasarathy, Phys.Rev. **D51**,5830 (1995).
- [7] T.L.Curtright, G.I.Ghandour and C.K.Zachos, Phys.Rev. **D34**, 3811 (1986).
- [8] W.Helfrich, Z.Naturforsch. **C28**, 693 (1973).
- [9] L.Peliti and S.Leibler, Phys.Rev.Lett. **54**, 1690 (1985).
- [10] R.Arnowitz, S.Deser, and C.W.Misner, in Gravitation, ed. L.Witten, (1962), New York, Wiley and sons.
- [11] C.W.Misner, K.S.Thorne, and J.A.Wheeler, Gravitation; W.H.Freeman and Co. San Francisco (1971).
- [12] A.Abrahams, A.Anderson, Y.Choquet-Bruhat and J.W.York, Phys.Rev.Lett. **75**, 3377 (1995).
- [13] A.A.Abrahams and J.W.York, Preprint- gr-qc/9601031 (1996).
- [14] A.Hosoya and Ken-ichi Nakao, Class.Quantum Gravity,**7**,163 (1990).
- [15] A.Hosoya and Ken-ichi Nakao, Prog.Theor.Phys. **84**, 739 (1990).
- [16] K.S.Viswanathan and R.Parthasarathy, IMSc preprint (1996).

- [17] W.Krivan and H.Herald, Class.Quantum Gravity. **12**, 2297 (1995).
- [18] J.Isenberg, Class.Quantum Gravity, **12**, 2249 (1995).
- [19] P.S.Letelier, Phys.Rev. **D41**, 1323 (1990).
- [20] B.Boisseau and P.S.Letelier, Phys.Rev. **D46**, 1721 (1992).
- [21] B.Carter, Phys.Rev. **D48**, 4835 (1993), Class.Quantum Grav. **11**, 2677 (1994).
- [22] J.Guven, Phys.Rev. **D48**, 4604,5562 (1993).
- [23] A.L.Larsen and V.P.Frolov, Nucl.Phys. **B414**, 129 (1994).
- [24] R.Capovilla and J.Guven, Phys.Rev. **D51**, 6736 (1995).
- [25] E.S.Fradkin and A.A.Tseytlin, Phys.Lett.B **158**, 316,(1985).
- [26] L.P.Eisenhart, Riemannian Geometry.
Princeton University Press. (1926).
- [27] K.Yano and M.Kon, Structures on Manifolds.
World Scientific. (1984).
- [28] A.P.Lightman, W.H.Press, R.H.Price and S.A.Teukolsky.
Problem Book in Relativity and Gravitation
Princeton University Press, Princeton, New Jersey, (1975).
- [29] W.Blaschke, vorlesungen über Differential Geometrie III.
Springer, Berlin (1929).
- [30] J.H.White, Proc.Amer.Math.Soc. **38**, 162 (1973).
- [31] T.J.Willmore, Total curvature in Riemannian Geometry.
Ellis Horwood Ltd. England. (1982).
- [32] S.Pokorski, Gauge Field Theories.
Cambridge University Press. Cambridge, (1987).
- [33] T.J.Willmore and C.S.Jhaveri, Quart.J.Math.Oxford (2) **23**, 319 (1972).

- [34] A.L.Kholodenko, Clemson University Preprint (1992).
- [35] L. Alvarez-Gaume, D.Z.Freedman, and S.Mukhi, Ann.of Phys. **134** 85 (1981)